

# Chapter 3. Discrete Random Variables and Probability Distributions

---

Math 3670 Summer 2024

Georgia Institute of Technology

# Section 1.

## Random Variables

---

# Random Variables

## Definition

For a given sample space  $\mathcal{S}$  of some experiment, a **random variable (RV)** is any rule that associates a number with each outcome in  $\mathcal{S}$ .

In mathematical language, a random variable  $X$  is a function whose domain is the sample space  $\mathcal{S}$  and whose range is the set of real numbers  $\mathbb{R}$ , that is

$$X : \mathcal{S} \rightarrow \mathbb{R}.$$

## Example

When a student calls a university help desk for technical support, he/she will either immediately be able to speak to someone ( $S$ , for success) or will be placed on hold ( $F$ , for failure).

With  $\mathcal{S} = \{S, F\}$ , define an RV  $X$  by

$$X(S) = 1, \quad X(F) = 0.$$

## Definition

Any random variable whose only possible values are 0 and 1 is called a **Bernoulli random variable**.

# Examples

## Example

Roll two dice.

Then,  $\mathcal{S} =$

Let

$X =$  The sum of two outcomes

$Y =$  The product of two outcomes

$U =$  The maximum of two outcomes

What are the possible values for  $X, Y, U$ ?

# Examples

## Example

Consider an experiment in which 9-volt batteries are tested until one with an acceptable voltage is obtained.

The sample space is  $\mathcal{S} = \{S, FS, FFS, \dots\}$ .

Let

$X =$  the number of batteries tested before the experiment terminates.

# Discrete Random Variables

## Definition

A **discrete random variable** is an RV whose possible values either constitute a finite set or else can be listed in an infinite sequence in which there is a first element, a second element, and so on (“countably” infinite).

Note: We will discuss **Continuous Random variables** later.



## Exercise

(3.1-4) Let  $X$  be the number of nonzero digits in a randomly selected zip code.

What are the possible values of  $X$ ?

Give three possible outcomes and their associated  $X$  values.

Section 2.  
Probability Distributions for  
Discrete Random Variables

---

## Definition

The probability mass function (PMF) of a discrete RV  $X$  is defined for every number  $x$  by

$$p(x) = \mathbb{P}(X = x) = \mathbb{P}(\{s \in \mathcal{S} : X(s) = x\}).$$

## Example

Consider whether the next person buying a computer at a certain electronics store buys a laptop or a desktop model. Let

$$X = \begin{cases} 1, & \text{if the customer purchases a desktop computer} \\ 0, & \text{if the customer purchases a laptop computer.} \end{cases}$$

If 20% of all purchasers during that week select a desktop, the PMF for  $X$  is

# Probability Mass Functions

## Example

Consider selecting at random a student who is among the 15,000 registered for the current term at Mega University.

Let  $X$  be the number of courses for which the selected student is registered with the PMF

$x$	1	2	3	4	5	6	7
$p(x)$	.01	.03	.13	.25	.39	.17	.02

What is the probability that the chosen student is registered for more than 4 courses?

# A Parameter of a Probability Distribution

## Example

Consider whether the next person buying a computer at a certain electronics store buys a laptop or a desktop model. Let

$$X = \begin{cases} 1, & \text{if the customer purchases a desktop computer} \\ 0, & \text{if the customer purchases a laptop computer.} \end{cases}$$

Then,  $X$  is a Bernoulli random variable.

If  $100\alpha\%$  of all purchasers during that week select a desktop, the PMF for  $X$  is

# A Parameter of a Probability Distribution

## Definition

Suppose  $p(x)$  depends on a quantity that can be assigned any one of a number of possible values, with each different value determining a different probability distribution.

Such a quantity is called a **parameter of the distribution**.

The collection of all probability distributions for different values of the parameter is called a **family of probability distributions**.

# A Parameter of a Probability Distribution

## Example

Starting at a fixed time, we observe the gender of each newborn child at a certain hospital until a boy (B) is born.

Let  $p = \mathbb{P}(B)$ , assume that successive births are independent, and define the RV  $X$  by the number of births observed.

Find the PMF of  $X$ .



# A Parameter of a Probability Distribution

## Definition

We say  $X$  is a **geometric random variable with success probability  $p$**  if its PMF is

$$p(x) =$$

# The Cumulative Distribution Function

## Definition

The cumulative distribution function (CDF)  $F(x)$  of a discrete RV variable  $X$  with PMF  $p(x)$  is defined by

$$F(x) =$$

# The Cumulative Distribution Function

## Example

Starting at a fixed time, we observe the gender of each newborn child at a certain hospital until a boy (B) is born.

Let  $p = \mathbb{P}(B)$ , assume that successive births are independent, and define the RV  $X$  by the number of births observed.

Find the CDF.

## Exercise

(3.2-17) A new battery's voltage may be acceptable (A) or unacceptable (U).

A certain flashlight requires two batteries, so batteries will be independently selected and tested until two acceptable ones have been found.

Suppose that 90% of all batteries have acceptable voltages.

Let  $Y$  denote the number of batteries that must be tested.

Find  $p(2)$ ,  $p(3)$ , and  $p(5)$ .

## Section 3.

### Expected Values

---

## Expected Values

Consider a university having 15,000 students and let  $X$  be the number of courses for which a randomly selected student is registered.

$x$	1	2	3	4	5	6	7
$p(x)$	.01	.03	.13	.25	.39	.17	.02
number registered	150	450	1950	3750	5850	2550	300

The average number of courses per student is

# Expected Values

## Definition

Let  $X$  be a discrete RV with set of possible values  $D$  and PMF  $p(x)$ .

The **expected value or mean value** of  $X$  is

$$\mathbb{E}[X] = \mu_X = \mu =$$

## Example

Let  $X$  be a Bernoulli random variable with  $p$ .

Find the expectation of  $X$ .



## Example

Let  $X$  be a geometric random variable with  $p$ .

Find the expectation of  $X$ .

# The Expected Value of a Function

## Example

Suppose a bookstore purchases ten copies of a book at \$6.00 each to sell at \$12.00 with the understanding that at the end of a 3-month period any unsold copies can be redeemed for \$2.00.

If  $X$  is the number of copies sold, then net revenue is

$$h(X) =$$

What then is the expected net revenue?

# The Expected Value of a Function

## Proposition

If the RV  $X$  has a set of possible values  $D$  and PMF  $p(x)$ , then the expected value of any function  $h(X)$  is

$$\mathbb{E}[h(X)] =$$

## Exercise

A computer store has purchased three computers of a certain type at \$500 apiece.

It will sell them for \$1000 apiece.

The manufacturer has agreed to repurchase any computers still unsold after a specified period at \$200 apiece.

Let  $X$  denote the number of computers sold with PMF

$$p(0) = .1, \quad p(1) = .2, \quad p(2) = .3, \quad p(3) = .4.$$

Find the expected profit

# Variance and Standard Deviation

## Definition

Let  $X$  have PMF  $p(x)$  and expected value  $\mu$ .

Then the variance of  $X$  is defined by

$$\text{Var}(X) = \sigma^2 = \sigma_X^2 =$$

The standard deviation (SD) of  $X$  is

$$\sigma_X = \sqrt{\text{Var}(X)}$$

# Variance and Standard Deviation

## Example

Let  $X$  be a discrete RV with PMF

$$p(0) = .1, \quad p(1) = .2, \quad p(2) = .3, \quad p(3) = .4.$$

Find the variance.

Proposition

$$\text{Var}(X) =$$

# Variance and Standard Deviation

## Example

Let  $X$  be a discrete RV with PMF

$$p(0) = .1, \quad p(1) = .2, \quad p(2) = .3, \quad p(3) = .4.$$

Find the variance using the formula.



## Proposition

$$\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$$
$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

## Exercise

(3.3-33) Let  $X$  be a Bernoulli RV with  $p$ .

Find the variance.

Section 4.  
The Binomial Probability  
Distribution

---

## Definition

An experiment consisting of a sequence of  $n$  trials is called a **Binomial experiment** if

1. Each trial can result in one of the same **two possible outcomes** (dichotomous trials), which we generically denote by success (S) and failure (F).
2. The trials are **independent**, so that the outcome on any particular trial does not influence the outcome on any other trial.
3. The probability of success  $\mathbb{P}(S)$  is **constant** from trial to trial; we denote this probability by  $p$ .

## Example

In a box, there are 7 Green balls and 3 Red balls.

Consider an experiment with 3 trials: we draw 3 balls in a row **with replacement**.

Let  $A_i$  be the event that the  $i$ -th ball is Green for  $i = 1, 2, 3$ .

1. Find  $\mathbb{P}(A_1), \mathbb{P}(A_2), \mathbb{P}(A_3)$ .
2. Are  $A_1, A_2$  independent?
3. Is this experiment binomial?

## Example

In a box, there are 7 Green balls and 3 Red balls.

Consider an experiment with 3 trials: we draw 3 balls in a row **without replacement**.

Let  $A_i$  be the event that the  $i$ -th ball is Green for  $i = 1, 2, 3$ .

1. Find  $\mathbb{P}(A_1), \mathbb{P}(A_2), \mathbb{P}(A_3)$ .
2. Are  $A_1, A_2$  independent?
3. Is this experiment binomial?

## Rule

Consider sampling without replacement from a dichotomous population of size  $N$ . If the sample size (number of trials)  $n$  is at most 5% of the population size, the experiment can be analyzed as though it were exactly a binomial experiment.

# The Binomial Random Variable and Distribution

## Definition

The **binomial random variable**  $X$  associated with a binomial experiment consisting of  $n$  trials is defined as **the number of Success among the  $n$  trials**.

We denote by  $X \sim \text{Bin}(n, p)$ .



# The Binomial Random Variable and Distribution

## Example

Consider tossing a coin 5 times.

Assume that the probability of landing Heads is 0.4.

What is the probability having Heads 3 times?

# The Binomial Random Variable and Distribution

## The PMF and CDF of Binomial RV

Let  $X \sim \text{Bin}(n, p)$ .

The PMF is

$$b(x; n, p) = \mathbb{P}(X = x) =$$

The CDF is

$$B(x; n, p) = \mathbb{P}(X \leq x) =$$

# The Binomial Random Variable and Distribution

## Example

Suppose that 20% of all copies of a particular textbook fail a certain binding strength test.

Let  $X$  denote the number among 15 randomly selected copies that fail the test.

1. How is  $X$  distributed? under what assumptions?
2. Find the probability that at most 8 fail the test.
3. Find the probability that exactly 8 fail the test.

# The Binomial Random Variable and Distribution

c.  $n = 15$

		<i>p</i>														
		0.01	0.05	0.10	0.20	0.25	0.30	0.40	0.50	0.60	0.70	0.75	0.80	0.90	0.95	0.99
	0	.860	.463	.206	.035	.013	.005	.000	.000	.000	.000	.000	.000	.000	.000	.000
	1	.990	.829	.549	.167	.080	.035	.005	.000	.000	.000	.000	.000	.000	.000	.000
	2	1.000	.964	.816	.398	.236	.127	.027	.004	.000	.000	.000	.000	.000	.000	.000
	3	1.000	.995	.944	.648	.461	.297	.091	.018	.002	.000	.000	.000	.000	.000	.000
	4	1.000	.999	.987	.836	.686	.515	.217	.059	.009	.001	.000	.000	.000	.000	.000
	5	1.000	1.000	.998	.939	.852	.722	.403	.151	.034	.004	.001	.000	.000	.000	.000
	6	1.000	1.000	1.000	.982	.943	.869	.610	.304	.095	.015	.004	.001	.000	.000	.000
<i>x</i>	7	1.000	1.000	1.000	.996	.983	.950	.787	.500	.213	.050	.017	.004	.000	.000	.000
	8	1.000	1.000	1.000	.999	.996	.985	.905	.696	.390	.131	.057	.018	.000	.000	.000
	9	1.000	1.000	1.000	1.000	.999	.996	.966	.849	.597	.278	.148	.061	.002	.000	.000
	10	1.000	1.000	1.000	1.000	1.000	.999	.991	.941	.783	.485	.314	.164	.013	.001	.000
	11	1.000	1.000	1.000	1.000	1.000	1.000	.998	.982	.909	.703	.539	.352	.056	.005	.000
	12	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.996	.973	.873	.764	.602	.184	.036	.000
	13	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.995	.965	.920	.833	.451	.171	.010
	14	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	.995	.987	.965	.794	.537	.140

# The Binomial Random Variable and Distribution

## Example

An electronics manufacturer claims that at most 10% of its power supply units need service during the warranty period.

To investigate this claim, technicians at a testing laboratory purchase 20 units. Let  $p$  denote the probability that a power supply unit needs repair. The laboratory technicians must decide whether the data resulting from the experiment supports the claim that  $p \leq .10$ .

Let  $X$  denote the number among the 20 sampled that need repair.

Reject the claim that  $p \leq .10$  in favor of the conclusion that  $p > .10$  if  $X \geq 5$ .

Probability of wrong conclusion?

# The Binomial Random Variable and Distribution

## Proposition

The mean and the variance of  $X \sim \text{Bin}(n, p)$  are

$$\mathbb{E}[X] =$$

$$\text{Var}(X) =$$

## Exercise

(3.4-52) Suppose that 30% of all students who have to buy a text for a particular course want a new copy (the successes!), whereas the other 70% want a used copy. Consider randomly selecting 25 purchasers.

1. What are the mean value and standard deviation of the number who want a new copy of the book?
2. What is the probability that the number who want new copies is more than two standard deviations away from the mean value?
3. Suppose that new copies cost \$100 and used copies cost \$70. Assume the bookstore currently has 50 new copies and 50 used copies. What is the expected value of total revenue from the sale of the next 25 copies purchased?

Section 5.  
Hypergeometric and Negative  
Binomial Distributions

---



# The Hypergeometric Distribution

## Definition

1. The population or set to be sampled consists of  $N$  individuals, objects, or elements (a finite population).
2. Each individual can be characterized as a success (S) or a failure (F), and there are  $M$  successes in the population.
3. A sample of  $n$  individuals is selected **without replacement** in such a way that each subset of size  $n$  is equally likely to be chosen.

Let  $X$  be the number of Success in the sample.

The random variable  $X$  is called **hypergeometric** and denoted by  $X \sim \text{HG}(n, M, N)$

# The Hypergeometric Distribution

## Example

During a particular period a university's information technology office received 20 service orders for problems with printers, of which 8 were laser printers and 12 were inkjet models.

Suppose that the 5 are selected in a completely random fashion, so that any particular subset of size 5 has the same chance of being selected as does any other subset.

What then is the probability that exactly 3 of the selected service orders were for inkjet printers?

# The Hypergeometric Distribution

## Proposition

Let  $X \sim \text{HG}(n, M, N)$ .

The PMF is

$$h(x; n, M, N) = \mathbb{P}(X = x) =$$

for  $x$  satisfying

# The Hypergeometric Distribution

## Example

Five individuals from an animal population thought to be near extinction in a certain region have been caught, tagged, and released to mix into the population.

After they have had an opportunity to mix, a random sample of 10 of these animals is selected.

Let  $X$  be the number of tagged animals in the second sample.

If there are actually 25 animals of this type in the region, what is the probability that (a)  $X = 2$  (b)  $X \leq 2$ ?

# The Hypergeometric Distribution

## Proposition

The mean and the variance of  $X \sim \text{HG}(n, M, N)$  are

$$\mathbb{E}[X] = n \cdot \frac{M}{N}$$
$$\text{Var}(X) = \left( \frac{N-n}{N-1} \right) \cdot n \cdot \frac{M}{N} \left( 1 - \frac{M}{N} \right)$$

# The Negative Binomial Distribution

## Definition

1. The experiment consists of a sequence of independent trials.
2. Each trial can result in either a success (S) or a failure (F).
3. The probability of success is constant from trial to trial.
4. The experiment continues (trials are performed) until a total of  $r$  successes have been observed, where  $r$  is a specified positive integer.

Let  $X$  be the number of failures that precede the  $r$ -th success.

The random variable  $X$  is called **Negative Binomial** and denoted by  $X \sim \text{NegBin}(r, p)$ .

# The Negative Binomial Distribution

## Proposition

The PMF of  $X \sim \text{NegBin}(r, p)$  is

$$nb(x; r, p) = \mathbb{P}(X = x) =$$

# The Negative Binomial Distribution

## Example

A pediatrician wishes to recruit 5 couples, each of whom is expecting their first child, to participate in a new natural childbirth regimen. Let

$$p = \mathbb{P}(\text{a randomly selected couple agrees to participate}).$$

If  $p = .2$ , what is the probability that 15 couples must be asked before 5 are found who agree to participate?



# The Negative Binomial Distribution

## Proposition

The mean and the variance of  $X \sim \text{NegBin}(r, p)$  are

$$\mathbb{E}[X] =$$

$$\text{Var}(X) =$$

## Exercise

(3.5-71) A geologist has collected 10 specimens of basaltic rock and 10 specimens of granite. The geologist instructs a laboratory assistant to randomly select 15 of the specimens for analysis.

1. What is the PMF of the number of granite specimens selected for analysis?
2. What is the probability that all specimens of one of the two types of rock are selected for analysis?
3. What is the probability that the number of granite specimens selected for analysis is within 1 standard deviation of its mean value?



Section 6.  
The Poisson Probability  
Distribution

---

## Definition

A discrete random variable  $X$  is said to have a Poisson distribution with parameter  $\mu > 0$  if the PMF of  $X$  is

$$p(x; \mu) = \mathbb{P}(X = x) =$$

We denote by  $X \sim \text{Pois}(\mu)$ .

1. The parameter  $\mu$  stands for
2. Recall that

$$e^x =$$

## Proposition

The mean and the variance of  $X \sim \text{Pois}(\mu)$  are

$$\mathbb{E}[X] =$$

$$\text{Var}(X) =$$

## Example

Let  $X$  denote the number of creatures of a particular type captured in a trap during a given time period.

Suppose that  $X$  has a Poisson distribution with  $\mu = 4.5$ , so on average traps will contain 4.5 creatures.

What is the probability that a trap contains exactly five creatures?

What are the expectation and the variance?

# The Poisson Distribution as a Limit

## Proposition

Suppose that in the binomial PMF  $b(x; n, p)$ , we let  $n \rightarrow \infty$  and  $p \rightarrow 0$  in such a way that  $np$  approaches a value  $\mu$ . Then

$$b(x; n, p) \rightarrow p(x; \mu).$$



# The Poisson Distribution as a Limit

According to this proposition, in any binomial experiment in which  $n$  is large and  $p$  is small,

$$b(x; n, p) \approx p(x; \mu),$$

where  $\mu = np$ .

As a rule of thumb, this approximation can safely be applied if  $n > 50$  and  $np < 5$ .

## Example

If a publisher of nontechnical books takes great pains to ensure that its books are free of typographical errors, so that the probability of any given page containing at least one such error is .005 and errors are independent from page to page,

what is the probability that one of its 400-page novels will contain exactly one page with errors? At most three pages with errors?

# The Poisson Process

Events of interest might be visits to a particular website, pulses of some sort recorded by a counter, email messages sent to a particular address, accidents in an industrial facility, or cosmic ray showers observed by astronomers at a particular observatory. We make the following assumptions about the way in which the events of interest occur:

1. There exists a parameter  $\alpha > 0$  such that for any short time interval of length  $\Delta t$ , the probability that exactly one event occurs is  $\alpha\Delta t + o(\Delta t)$ .
2. The probability of more than one event occurring during  $\Delta t$  is  $o(\Delta t)$ .
3. The number of events occurring during the time interval  $\Delta t$  is independent of the number that occur prior to this time interval.

## Proposition

The number of events during a time interval of length  $t$  is a Poisson RV with parameter  $\mu = \alpha t$ .

## Example

Suppose pulses arrive at a counter at an average rate of six per minute, so that  $\alpha = 6$ . Find the probability that in a .5-min interval at least one pulse is received.

## Exercise

(3.6-86) The number of people arriving for treatment at an emergency room can be modeled by a Poisson process with a rate parameter of five per hour.

1. What is the probability that exactly four arrivals occur during a particular hour?
2. What is the probability that at least four people arrive during a particular hour?
3. How many people do you expect to arrive during a 45-min period?

