

# Chapter 5. Joint Probability Distributions and Random Samples

---

Math 3670 Spring 2025

Georgia Institute of Technology

Section 1.  
Jointly Distributed Random  
Variables

---

# Two Discrete Random Variables

## Definition

Let  $X$  and  $Y$  be two discrete RVs defined on the sample space  $\mathcal{S}$  of an experiment.

The joint probability mass function  $p(x,y)$  is defined by

$$p(x,y) =$$

The joint PMF satisfies

1.  $p(x,y) \geq 0$
2.  $\sum_{x,y} p(x,y) = 1$
3.  $\mathbb{P}((X,Y) \in A) = \sum_{(x,y) \in A} p(x,y)$

# Two Discrete Random Variables

## Example

A large insurance agency services a number of customers who have purchased both a homeowner's policy and an automobile policy from the agency. For an automobile policy, the choices are \$100 and \$250, whereas for a homeowner's policy, the choices are 0, \$100, and \$200.

Suppose an individual with both types of policy is selected at random from the agency's files. Let  $X$  be deductible amount on the auto policy and  $Y$  deductible amount on the homeowner's policy.

	0	100	200
100	0.2	0.1	0.2
250	0.05	0.15	0.3

# Two Discrete Random Variables

## Definition

For a given joint PMF  $p(x,y)$  of random variables  $X$  and  $Y$ , the marginal probability mass function of  $X$  is given by

$$p_X(x) :=$$

# Two Discrete Random Variables

## Example

A large insurance agency services a number of customers who have purchased both a homeowner's policy and an automobile policy from the agency. For an automobile policy, the choices are \$100 and \$250, whereas for a homeowner's policy, the choices are 0, \$100, and \$200.

Suppose an individual with both types of policy is selected at random from the agency's files. Let  $X$  be deductible amount on the auto policy and  $Y$  deductible amount on the homeowner's policy.

	0	100	200
100	0.2	0.1	0.2
250	0.05	0.15	0.3

# Two Continuous Random Variables

## Definition

Let  $X$  and  $Y$  be two continuous RVs.

The joint probability density function  $f(x, y)$  is defined by

$$f(x, y) =$$

The joint PDF satisfies

1.  $f(x, y) \geq 0$
2.  $\int f(x, y) = 1$
3.  $\mathbb{P}((X, Y) \in A) = \int_A f(x, y)$

# Two Continuous Random Variables

## Example

A bank operates both a drive-up facility and a walk-up window.

On a randomly selected day, let  $X$  be the proportion of time that the drive-up facility is in use (at least one customer is being served or waiting to be served) and  $Y$  the proportion of time that the walk-up window is in use.

The joint PDF is given by

$$f(x, y) = \begin{cases} \frac{6}{5}(x + y^2), & 0 \leq x \leq 1, 0 \leq y \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Find  $\mathbb{P}(0 \leq X \leq \frac{1}{4}, 0 \leq Y \leq \frac{1}{4})$ .



# Two Continuous Random Variables

## Definition

For a given joint PDF  $f(x,y)$  of random variables  $X$  and  $Y$ , the marginal probability density function of  $X$  is given by

$$f_X(x) :=$$

# Two Continuous Random Variables

## Example

A bank operates both a drive-up facility and a walk-up window.

On a randomly selected day, let  $X$  be the proportion of time that the drive-up facility is in use (at least one customer is being served or waiting to be served) and  $Y$  the proportion of time that the walk-up window is in use.

The joint PDF is given by

$$f(x,y) = \begin{cases} \frac{6}{5}(x + y^2), & 0 \leq x, y \leq 1 \\ 0 & \text{otherwise.} \end{cases}$$

Find the marginal PDFs.

# Two Continuous Random Variables

## Example

The joint PDF is given by

$$f(x,y) = \begin{cases} 24xy, & 0 \leq x \leq 1, 0 \leq y \leq 1, x + y \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Find the marginal PDFs and  $\mathbb{P}(X + Y \leq 1/2)$ .

# Independent Random Variables

## Definition

Two random variables  $X$  and  $Y$  are said to be **independent** if

# Independent Random Variables

## Example

The joint PDF is given by

$$f(x, y) = \begin{cases} 24xy, & 0 \leq x \leq 1, 0 \leq y \leq 1, x + y \leq 1 \\ 0 & \text{otherwise.} \end{cases}$$

Are X and Y independent?

# Independent Random Variables

## Example

Suppose that the lifetimes of two components are independent of one another and that the first lifetime,  $X_1$ , has an exponential distribution with parameter  $\lambda_1$ , whereas the second,  $X_2$ , has an exponential distribution with parameter  $\lambda_2$ .

Find the joint PDF.

# Independent Random Variables

## Definition

The random variables  $X_1, X_2, \dots, X_n$  are said to be **independent** if

# Conditional Distributions

## Definition

Let  $X$  and  $Y$  be two continuous RVs with joint PDF  $f(x, y)$ .

Then for any  $x$  for which  $f_X(x) > 0$ , **the conditional probability density function** of  $Y$  given that  $X = x$  is



# Conditional Distributions

## Example

A bank operates both a drive-up facility and a walk-up window.

On a randomly selected day, let  $X$  be the proportion of time that the drive-up facility is in use (at least one customer is being served or waiting to be served) and  $Y$  the proportion of time that the walk-up window is in use.

The joint PDF is given by

$$f(x, y) = \begin{cases} \frac{6}{5}(x + y^2), & 0 \leq x \leq 1, 0 \leq y \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Find the conditional PDF of  $Y$  given  $X = 0.8$ .

Compute  $\mathbb{P}(Y \leq 0.5 | X = 0.8)$ .

## Exercise

(5.1-12) Two components of a minicomputer have the following joint PDF for their useful lifetimes  $X$  and  $Y$ :

$$f(x, y) = \begin{cases} xe^{-x(y+1)}, & x \geq 0, y \geq 0, \\ 0 & \text{otherwise.} \end{cases}$$

1. What is the probability that the lifetime  $X$  of the first component exceeds 3?
2. What are the marginal PDFs of  $X$  and  $Y$ ? Are the two lifetimes independent? Explain.
3. What is the probability that the lifetime of at least one component exceeds 3?

## Section 2. Expected Values, Covariance, and Correlation

---

# Expectation of a Function of Two Random Variables

## Proposition

Let  $X$  and  $Y$  be jointly distributed RVs with PMF  $p(x, y)$  or PDF  $f(x, y)$  according to whether the variables are discrete or continuous.

Let  $h(x, y)$  be a function of two variables, then we can define a new random variable  $Z = h(X, Y)$ .

The expectation of  $Z$  is

$$\mathbb{E}[Z] = \mathbb{E}[h(X, Y)] =$$

# Expectation of a Function of Two Random Variables

## Example

Five friends have purchased tickets to a certain concert.

If the tickets are for seats 1–5 in a particular row and the tickets are randomly distributed among the five, what is the expected number of seats separating any particular two of the five?

Let  $X$  and  $Y$  denote the seat numbers of the first and second individuals, respectively.

# Expectation of a Function of Two Random Variables

## Example

The joint PDF is given by

$$f(x, y) = \begin{cases} 24xy, & 0 \leq x \leq 1, 0 \leq y \leq 1, x + y \leq 1 \\ 0 & \text{otherwise.} \end{cases}$$

Find  $\mathbb{E}[XY]$ .

## Definition

The covariance between two RVs  $X$  and  $Y$  is

$$\text{Cov}(X, Y) =$$

## Example

The joint PDF is given by

$$f(x, y) = \begin{cases} 24xy, & 0 \leq x \leq 1, 0 \leq y \leq 1, x + y \leq 1 \\ 0 & \text{otherwise.} \end{cases}$$

Find the covariance of  $X, Y$ .



Proposition

$$\text{Cov}(X, Y) =$$

## Definition

The correlation coefficient of  $X$  and  $Y$  is defined by

$$\text{Corr}(X, Y) = \rho_{X,Y} = \rho =$$

## Properties

1. For constants  $a, b, c, d$ ,

$$\text{Corr}(aX + b, cX + d) =$$

2.  $-1 \leq \text{Corr}(X, Y) \leq 1$
3. If  $X$  and  $Y$  are independent, then  $\text{Corr}(X, Y) = 0$ . The converse does not hold in general.
4. If  $\text{Corr}(X, Y) = 1, -1$ , then  $Y = aX + b$  for some  $a, b$ .

# Correlation Coefficient

## Example

The joint PMF is given by

$$p(x,y) = \begin{cases} \frac{1}{4}, & (x,y) = (-4, 1), (4, -1), (2, 2), (-2, -2) \\ 0 & \text{otherwise.} \end{cases}$$

Find the covariance and the correlation coefficient.

## Exercise

(5.2-24) Six individuals, including A and B, take seats around a circular table in a completely random fashion.

Suppose the seats are numbered  $1, 2, \dots, 6$ .

Let  $X$  be A's seat number and  $Y$  B's seat number.

If A sends a written message around the table to B in the direction in which they are closest,

how many individuals (including A and B) would you expect to handle the message?

Section 4.

# The Distribution of the Sample Mean

---

# Sample Mean

## Definition

The RVs  $X_1, X_2, \dots, X_n$  are said to form a (simple) random sample of size  $n$  if

1. they are independent RVs, and
2. every  $X_i$  has the same probability distribution.

The sample mean is defined by

$$\bar{X} =$$

The sample total is defined by

$$T =$$

## Proposition

Let  $X_1, \dots, X_n$  be a random sample from a distribution with mean  $\mu$  and variance  $\sigma^2$ .  
Then,

$$\mathbb{E}[\bar{X}] =$$

$$\text{Var}(\bar{X}) =$$

$$\mathbb{E}[T] =$$

$$\text{Var}(T) =$$



## Example

In a notched tensile fatigue test on a titanium specimen, the expected number of cycles to first acoustic emission (used to indicate crack initiation) is  $\mu = 28,000$ , and the standard deviation of the number of cycles is  $\sigma = 5000$ .

Let  $X_1, X_2, \dots, X_{25}$  be a random sample of size 25, where each  $X_i$  is the number of cycles on a different randomly selected specimen.

# The Case of a Normal Population Distribution


## Proposition

Let  $X_1, \dots, X_n$  be a random sample from a normal distribution with mean  $\mu$  and variance  $\sigma^2$ . Then,

$$\bar{X} \sim$$

$$T \sim$$

# The Case of a Normal Population Distribution



**Figure 5.14** A normal population distribution and  $\bar{X}$  sampling distributions

# The Case of a Normal Population Distribution

## Example

The time that it takes a randomly selected rat of a certain subspecies to find its way through a maze is a normally distributed RV with  $\mu = 1.5$  min and  $\sigma = .35$  min.

Suppose five rats are selected.

Let  $X_1, \dots, X_5$  denote their times in the maze.

Assuming the  $X_i$ 's to be a random sample from this normal distribution, what is the probability that the total time is between 6 and 8 min?

## Theorem

Let  $X_1, X_2, \dots, X_n$  be a random sample from a distribution with mean  $\mu$  and variance  $\sigma^2$ .

If  $n$  is sufficiently large,  $\bar{X}$  and  $T$  have approximately normal distributions.

Rule of Thumb: If  $n \geq 30$ , the Central Limit Theorem can be used.

## Example

A certain consumer organization customarily reports the number of major defects for each new automobile that it tests.

Suppose the number of such defects for a certain model is a random variable with mean value 3.2 and standard deviation 2.4.

Among 100 randomly selected cars of this model, how likely is it that the sample average number of major defects exceeds 4?

## Normal approximation to Binomial

If  $X \sim \text{Bin}(n, p)$  and  $n$  is large enough,

$X$  and  $X/n$  have approximately normal distribution.

## Exercise

(5.4-56) A binary communication channel transmits a sequence of “bits” (0s and 1s). Suppose that for any particular bit transmitted, there is a 10% chance of a transmission error (a 0 becoming a 1 or a 1 becoming a 0).

Assume that bit errors occur independently of one another.

1. Consider transmitting 1000 bits. What is the approximate probability that at most 125 transmission errors occur?
2. Suppose the same 1000-bit message is sent two different times independently of one another.

What is the approximate probability that the number of errors in the first transmission is within 50 of the number of errors in the second?



Section 5.  
The Distribution of a Linear  
Combination

---

# Linear Combination

## Definition

Given a collection of  $n$  random variables  $X_1, \dots, X_n$  and constants  $a_1, \dots, a_n$ ,

$$Y = a_1X_1 + a_2X_2 + \dots + a_nX_n$$

is called **a linear combination** of the  $X_i$ 's.

# Linear Combination

## Proposition

For a collection of  $n$  random variables  $X_1, \dots, X_n$  and constants  $a_1, \dots, a_n$ , consider

$$Y = a_1X_1 + a_2X_2 + \dots + a_nX_n$$

Then,

$$\mathbb{E}[Y] =$$

$$\text{Var}(Y) =$$

In particular, if they are independent,

$$\text{Var}(Y) =$$

## Example

A certain automobile manufacturer equips a particular model with either a six-cylinder engine or a four-cylinder engine.

Let  $X_1$  and  $X_2$  be fuel efficiencies for independently and randomly selected six-cylinder and four-cylinder cars, respectively, with

$$\mu_1 = 22, \quad \mu_2 = 26, \quad \sigma_1 = 1.2, \quad \sigma_2 = 1.5.$$

Find  $\mathbb{E}[X_1 - X_2]$  and  $\text{Var}(X_1 - X_2)$ .

## Proposition

If  $X_1, X_2, \dots, X_n$  are independent, normally distributed RVs (with possibly different means and/or variances), then any linear combination also has a normal distribution.

In particular, the difference  $X_1 - X_2$  between two independent, normally distributed variables is itself normally distributed.

## Exercise

(5.5-62) Manufacture of a certain component requires three different machining operations.

Machining time for each operation has a normal distribution, and the three times are independent of one another.

The mean values are 15, 30, and 20 min, respectively, and the standard deviations are 1, 2, and 1.5 min, respectively.

What is the probability that it takes at most 1 hour of machining time to produce a randomly selected component?