

# In-Class Final Exam Review Set A, Math 1554, Fall 2019

1. Indicate whether the statements are true or false.

true    false

- 
- |                       |                       |  |
|-----------------------|-----------------------|--|
| <input type="radio"/> | <input type="radio"/> | If a linear system has more unknowns than equations, then the system has either no solutions or infinitely many solutions. |
| <input type="radio"/> | <input type="radio"/> | A $n \times n$ matrix $A$ and its echelon form $E$ will always have the same eigenvalues.                                  |
| <input type="radio"/> | <input type="radio"/> | $x^2 - 2xy + 4y^2 \geq 0$ for all real values of $x$ and $y$ .   |
| <input type="radio"/> | <input type="radio"/> | If matrix $A$ has linearly dependent columns, then $\dim((\text{Row } A)^\perp) > 0$ .                                     |
| <input type="radio"/> | <input type="radio"/> | If $\lambda$ is an eigenvalue of $A$ , then $\dim(\text{Null}(A - \lambda I)) > 0$ .                                       |
| <input type="radio"/> | <input type="radio"/> | If $A$ has $QR$ decomposition $A = QR$ , then $\text{Col } A = \text{Col } Q$ .  |
| <input type="radio"/> | <input type="radio"/> | If $A$ has $LU$ decomposition $A = LU$ , then $\text{rank}(A) = \text{rank}(U)$ .  |
| <input type="radio"/> | <input type="radio"/> | If $A$ has $LU$ decomposition $A = LU$ , then $\dim(\text{Null } A) = \dim(\text{Null } U)$ .                              |
- 

2. Give an example of the following.

i) A  $4 \times 3$  lower triangular matrix,  $A$ , such that  $\text{Col}(A)^\perp$  is spanned by


the vector  $\vec{v} = \begin{pmatrix} 1 \\ 2 \\ 3 \\ 1 \end{pmatrix}$ .     $A = \begin{pmatrix} & & & \\ & & & \\ & & & \\ & & & \end{pmatrix}$

ii) A  $3 \times 4$  matrix  $A$ , that is in RREF, and satisfies  $\dim((\text{Row } A)^\perp) = 2$  and  $\dim((\text{Col } A)^\perp) =$


2.  $A = \begin{pmatrix} & & & \\ & & & \\ & & & \end{pmatrix}$

3. (3 points) Suppose  $A = \begin{pmatrix} 3 & 1 \\ 6 & 2 \end{pmatrix}$ . On the grid below, sketch a)  $\text{Col}(A)$ , and b) the eigenspace corresponding to eigenvalue  $\lambda = 5$ .

(a)  $\text{Col}(A)$



(b)  $\lambda = 5$  eigenspace



4. Fill in the blanks.

(a) If  $A \in \mathbb{R}^{M \times N}$ ,  $M < N$ , and  $A\vec{x} = 0$  does not have a non-trivial solution, how many pivot columns does  $A$  have?

(b) Consider the following linear transformation.

$$T(x_1, x_2) = (2x_1 - x_2, 4x_1 - 2x_2, x_2 - 2x_1).$$

The domain of  $T$  is . The image of  $\vec{x} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$  under  $T(\vec{x})$  is  $\begin{pmatrix} \phantom{0} \\ \phantom{0} \\ \phantom{0} \end{pmatrix}$ . The co-domain of  $T$  is . The range of  $T$  is:

5. Four points in  $\mathbb{R}^2$  with coordinates  $(t, y)$  are  $(0, 1)$ ,  $(\frac{1}{4}, \frac{1}{2})$ ,  $(\frac{1}{2}, -\frac{1}{2})$ , and  $(\frac{3}{4}, -\frac{1}{2})$ . Determine the values of  $c_1$  and  $c_2$  for the curve  $y = c_1 \cos(2\pi t) + c_2 \sin(2\pi t)$  that best fits the points. Write the values you obtain for  $c_1$  and  $c_2$  in the boxes below.

$$c_1 = \text{} \quad c_2 = \text{$$

## In-Class Final Exam Review Set B, Math 1554, Fall 2019

1. Indicate whether the statements are true or false.

true    false

---

- |                       |                       |   |
|-----------------------|-----------------------|---|
| <input type="radio"/> | <input type="radio"/> | For any vector $\vec{y} \in \mathbb{R}^2$ and subspace $W$ , the vector $\vec{v} = \vec{y} - \text{proj}_W \vec{y}$ is orthogonal to $W$ .            |
| <input type="radio"/> | <input type="radio"/> | If $A$ is $m \times n$ and has linearly dependent columns, then the columns of $A$ cannot span $\mathbb{R}^m$ .                                       |
| <input type="radio"/> | <input type="radio"/> | If a matrix is invertible it is also diagonalizable.  |
| <input type="radio"/> | <input type="radio"/> | If $E$ is an echelon form of $A$ , then $\text{Null } A = \text{Null } E$ .   |
| <input type="radio"/> | <input type="radio"/> | If the SVD of $n \times n$ singular matrix $A$ is $A = U\Sigma V^T$ , then $\text{Col}A = \text{Col}U$ .  |
| <input type="radio"/> | <input type="radio"/> | If the SVD of $n \times n$ matrix $A$ is $A = U\Sigma V^T$ , $r = \text{rank}A$ , then the first $r$ columns of $V$ give a basis for $\text{Null}A$ . |
- 

2. Give an example of:


a) a vector  $\vec{u} \in \mathbb{R}^3$  such that  $\text{proj}_{\vec{p}} \vec{u} = \vec{p}$ , where  $\vec{u} \neq \vec{p}$ , and  $\vec{p} = \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix}$ :  $\vec{u} = \begin{pmatrix} \phantom{0} \\ \phantom{2} \\ \phantom{0} \end{pmatrix}$

b) an upper triangular  $4 \times 4$  matrix  $A$  that is in RREF, 0 is its only eigenvalue, and its corresponding eigenspace is 1-dimensional.  $A = \begin{pmatrix} \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} \end{pmatrix}$

c) A  $3 \times 4$  matrix,  $A$ , and  $\text{Col}(A)^\perp$  is spanned by  $\begin{pmatrix} 1 \\ -3 \\ -4 \end{pmatrix}$ .

d) A  $2 \times 2$  matrix in RREF that is diagonalizable and not invertible.

3. Suppose  $A = \begin{pmatrix} 2 & -1 \\ 4 & -2 \end{pmatrix}$ . On the grid below, sketch a) the range of  $x \rightarrow Ax$ , b)  $(\text{Col } A)^\perp$ , (c) set of solutions to  $A\vec{x} = \begin{pmatrix} 3 \\ 6 \end{pmatrix}$ .



4. Matrix  $A$  is a  $2 \times 2$  matrix whose eigenvalues are  $\lambda_1 = \frac{1}{2}$  and  $\lambda_2 = 1$ , and whose corresponding eigenvectors are  $\vec{v}_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ ,  $\vec{v}_2 = \begin{pmatrix} 4 \\ 1 \end{pmatrix}$ . Calculate

1.  $A(\vec{v}_1 + 4\vec{v}_2)$
2.  $A^{10}$
3.  $\lim_{k \rightarrow \infty} A^k(\vec{v}_1 + 4\vec{v}_2)$

## In-Class Final Exam Review Set C, Math 1554, Fall 2019

1. Indicate whether the statements are possible or impossible.

possible    impossible

---

- |                       |                       |   |
|-----------------------|-----------------------|---|
| <input type="radio"/> | <input type="radio"/> | $Q(\vec{x}) = \vec{x}^T A \vec{x}$ is a positive definite quadratic form, and $Q(\vec{v}) = 0$ , where $\vec{v}$ is an eigenvector of $A$ .                                       |
| <input type="radio"/> | <input type="radio"/> | The maximum value of $Q(\vec{x}) = ax_1^2 + bx_2^2 + cx_3^2$ , where $a > b > c$ , for $\vec{x} \in \mathbb{R}^3$ , subject to $\ \vec{x}\  = 1$ , is not unique.                 |
| <input type="radio"/> | <input type="radio"/> | The location of the maximum value of $Q(\vec{x}) = ax_1^2 + bx_2^2 + cx_3^2$ , where $a > b > c$ , for $\vec{x} \in \mathbb{R}^3$ , subject to $\ \vec{x}\  = 1$ , is not unique. |
| <input type="radio"/> | <input type="radio"/> | $A$ is $2 \times 2$ , the algebraic multiplicity of eigenvalue $\lambda = 0$ is 1, and $\dim(\text{Col}(A)^\perp)$ is equal to 0.   |
| <input type="radio"/> | <input type="radio"/> | Stochastic matrix $P$ has zero entries and is regular.  |
| <input type="radio"/> | <input type="radio"/> | $A$ is a square matrix that is not diagonalizable, but $A^2$ is diagonalizable.   |
| <input type="radio"/> | <input type="radio"/> | The map $T_A(\vec{x}) = A\vec{x}$ is one-to-one but not onto, $A$ is $m \times n$ , and $m < n$ .   |
- 

2. Transform  $T_A = A\vec{x}$  reflects points in  $\mathbb{R}^2$  through the line  $y = 2 + x$ . Construct a standard matrix for the transform using homogeneous coordinates. Leave your answer as a product of three matrices.

3. Fill in the blanks.

- (a)  $T_A = A\vec{x}$ , where  $A \in \mathbb{R}^{2 \times 2}$ , is a linear transform that first rotates vectors in  $\mathbb{R}^2$  clockwise by  $\pi/2$  radians about the origin, then reflects them through the line  $x_1 = x_2$ . What is the value of  $\det(A)$ ?
- (b)  $B$  and  $C$  are square matrices with  $\det(BC) = -5$  and  $\det(C) = 2$ . What is the value of  $\det(B) \det(C^4)$ ?
- (c)  $A$  is a  $6 \times 4$  matrix in RREF, and  $\text{rank}(A) = 4$ . How many different matrices can you construct that meet these criteria?
- (d)  $T_A = A\vec{x}$ , where  $A \in \mathbb{R}^{2 \times 2}$ , projects points onto the line  $x_1 = x_2$ . What is an eigenvalue of  $A$  equal to?
- (e) If an eigenvalue of  $A$  is  $\frac{1}{3}$ , what is one eigenvalue of  $A^{-1}$  equal to?
- (f) If  $A$  is  $30 \times 12$  and  $A\vec{x} = \vec{b}$  has a unique least squares solution  $\hat{x}$  for every  $\vec{b}$  in  $\mathbb{R}^{30}$ , the dimension of  $\text{Null}A$  is .

4.  $A$  is a  $2 \times 2$  matrix whose nullspace is the line  $x_1 = x_2$ , and  $C = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$ . Sketch the nullspace of  $Y = AC$ .

5. Construct an SVD of  $A = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}$ . Use your SVD to calculate the condition number of  $A$ .